

# Pest News

A newsletter about pest management in Canterbury

## Wandering wallabies

Environment Canterbury would like to hear of any sightings of wallabies in North Canterbury. Reports have been received of wallabies on Banks Peninsula, at Mt Oxford and in other areas of Canterbury. Sightings outside of the South Canterbury containment zone suggest that animals are being illegally released. It is an offence to release a wallaby or to keep one as a pet.

Wallabies outside the containment zone pose a real threat to the pastoral economy and to native biodiversity. They feed mainly on grasses and herbs, cropping vegetation very close to the ground. This creates bare patches which then allow the soil to be eroded by wind and rain.

Exotic tree plantations are vulnerable to wallaby damage during the establishment stage and there is evidence of wallabies grazing on green feed crops, particularly when there is suitable cover nearby. Wallabies also prevent the regeneration of native bush by damaging the forest understorey when they are present in sufficient numbers.

One animal can eat the same amount as an average breeding ewe.

At low population densities, scat (droppings) and foot prints are good ways to detect the presence of wallabies.

If you suspect you may have wallabies on your property, looking for 'sign' is a good way to confirm their presence.

**Please contact your local Biosecurity Officer at Amberley to report wallaby sign or a sighting.**



Wallaby tracks are wide and flat. The hind foot leaves a print quite distinct from any other animal track you are likely to find in New Zealand.



Wallaby scat (droppings) is ovate in shape, dry and fibrous.

## Have you seen this plant?

Darwin's barberry (*Berberis darwinii*) is a spiny, evergreen shrub that can grow up to about 4 metres high. It can be found on forest and bush margins, sometimes in pasture and in ungrazed areas. Darwin's barberry has small clusters of glossy dark green leaves up to 3 centimetres long, with spiny edges. Small yellow-orange flowers are produced in clusters from September through to February. The flowers are followed by hanging groups of purple-black berries, which are a very popular food source for birds and consequently lead to the further spread of Darwin's barberry.

There are only a handful of known sites in Canterbury north of the Waimakariri River. Please inform Environment Canterbury's Biosecurity staff if you think you have seen this plant. Contact your local Biosecurity Officer at our Amberley office to report your sighting. Don't bring samples into the office as this risks spreading it further.



The dark green leaves and yellow-orange buds (left) and flowers (right) of Darwin's barberry

### Waikari Biosecurity Staff members

- Laurence Smith, Biosecurity Team Leader** .....Ph: 03 314 9035
- Leanne Lye, Biosecurity Support Officer** .....Ph: 03 314 8014
- Lindsay Scott, Biosecurity Officer** .....Ph: 03 314 9513
- Marika Fitt, Biosecurity Officer**.....Ph: 03 314 9582
- Lance Smith, Biosecurity Officer**.....Ph: 03 314 9585
- Terry Charles Biosecurity Officer (Rabbits)** .....Ph: 03 314 8403

### Waikari Pest Management Liaison Committee members

- Harry Pawsey (Chairperson)** ..... Ph 03 314 4169
- Ian Ferguson** ..... Ph 03 314 4133
- Nigel Fraser** ..... Ph 03 314 3423
- Robin Gardiner** ..... Ph 03 314 6744
- Kevin Earl** ..... Ph 03 314 4305
- Jock Foster**..... Ph 03 314 3831
- Graham Gibb** ..... Ph 03 314 3872
- Ben Cassidy** ..... Ph 03 314 4503
- Ted Phipps**..... Ph 03 314 4343
- Malcolm Gilbert** ..... Ph 03 314 5831
- Tim Cameron** ..... Ph 03 314 4575
- James Forbes** ..... Ph 03 314 6865
- Winton Dalley (HDC Rep)**..... Ph 03 314 4230

### A few words from Harry Pawsey, Pest Committee Chair

It has been a funny old year. The rainmaker holidayed from early January through to June, apologised for the distress it caused by raining until September. Then to add the final insult, presented a cold spring. This inclement weather has not deterred pests from multiplying, particularly rabbits. The best way to maximise the benefit of RHD is to kill the survivors of a virus outbreak.

Environment Canterbury, supported by Federated Farmers and the pest liaison committees, has applied for a global consent for the application of 1080 and Pindone throughout the Canterbury Region. If successful, this will remove the need for costly individual consents to apply these toxins. If you need assistance to coordinate a multi-property control program contact Steve Palmer, Airon Pest Solutions Ltd. on 03 319 8400 or 027 348 5394, his services are paid for by rates.

The Tb campaign continues to progress, with the number of Tb infected herds between the Waipara and Hurunui Rivers falling by one, to four.

Our vigilant biosecurity officers have responded quickly to sightings of wallabies around Leithfield. This is one pest we certainly don't need as they are very expensive to eradicate.

Many land occupiers are commenting on an increase of scattered broom and gorse on their own and neighbouring properties. Prudent land occupiers will eradicate scattered plants before they quickly become a major and very expensive problem.

Biosecurity staff continue control work on the Cheviot Chilean needle grass infestation at Spotswood. Their early action seems to have contained the infestation and they are still working on public awareness to ensure people are on the lookout for this plant.

My best wishes to all for the coming New Year, my wish being for a moist autumn for a change.

## CHILEAN NEEDLE GRASS *Nassella neesiana*



Chilean needle grass (*Nassella neesiana*) or CNG was first recorded in New Zealand in the 1930's. There are only two known areas of widespread infestation in the country - in Hawkes Bay and in Marlborough. The limited distribution is possibly due to the heavy seed produced by this plant which tends to fall close to the parent plant unless spread in hay, by stock, people or vehicle movement.

A routine inspection as part of Environment Canterbury's pest surveillance programme in November 2008 found the first incidence of Chilean needle grass in Canterbury. The plant was found growing at a vineyard at Spotswood just north of Cheviot. A thorough search of the area by Biosecurity staff found the infestation covered more 70 hectares with a handful of isolated patches nearby.

Difficult to identify, Chilean needle grass blends in well with other pasture grasses particularly in the spring growth period. CNG leaves are bright green, ribbed on the upper surface and rough to the touch. The plant becomes more distinctive during flowering and seeding when seedheads have a purplish tinge and the seed has a distinctive long twisting tail.

Chilean needle grass is unpalatable to stock when seeding and forms dense stands in pasture, dramatically reducing productivity after only a few years. The seed is extremely sharp and hairy and so catches in the pelt of passing animals. The seed tail has a corkscrew shape and so is difficult to remove. Like barley grass, the seed can work its way into tissue causing abscessing and downgrading hides and carcasses.

Chilean needle grass is extremely difficult to eradicate once established and so strict containment processes have been put in place at Spotswood to prevent spread from the area.

So far this season Biosecurity staff have grubbed and sprayed the main sites at Spotswood. Further searches will be carried out around the area over coming months and follow-up control work undertaken as needed.

**If you have found a plant on your property which you think may be Chilean needle grass please contact Environment Canterbury's Amberley Office on 03 314 8014. Do not bring in any samples for identification as this may increase the risk of spread.**

## Controlling gorse and broom



Gorse and broom on your property should be controlled when plants are actively growing as they will take up applied chemicals readily. Ideally, plants should be treated after flowering, but before seed set.

The priorities for your control programme should be:

- **Clear boundaries** of all gorse and broom at least 10m back (where those boundaries are clear or being cleared of gorse and broom)
- **Focus on clearing scattered plants and small patches.** These are the biggest threat to your property.
- **Plan and work methodically** across the property, beginning with the least infested areas and working back towards block infestations
- **Always control plants before they set seed** to prevent more seed being added to the seed bank
- **Trim boundary hedges** (top back and sides) each year after flowering but before they set seed.

If you would like assistance in developing a gorse and broom control programme for your property or would like to discuss your current control program, please contact a Biosecurity Officer at Environment Canterbury's Amberley Office for advice.

## Wilding conifer control update

Wilding conifers are trees that occur through wind-spread seed falling outside areas of managed trees. Lodgepole pine (*Pinus contorta*), Mountain pine (*P. mugo*), Corsican pine (*P. nigra*) and Douglas fir (*Pseudotsuga menziesii*) all occur as wilding conifers. In areas of Canterbury such as Craigieburn and Lake Pukaki, thousands of hectares of tussock grassland are infested with an almost blanket cover of wilding conifers of various ages.

Careful site selection for new conifer plantations is a must in order to prevent the spread of wilding conifers in future. Douglas fir in particular shows a clear tendency to take over clean tussock country. It is also of major concern because of its shade tolerance – it can germinate and grow under a native canopy and can out-compete native trees, making it a particular threat to biodiversity.

In North Canterbury between the Waimakariri River and Canterbury's northern boundary, wilding conifers have become established and are now present in large numbers, making control work a major undertaking.

Environment Canterbury maintains a close working relationship with the Department of Conservation and local land occupiers throughout Canterbury for the control of wilding conifers. Work in the Hurunui district is planned to complement the programmes of all concerned parties.



Environment Canterbury began control operations in the Hurunui River Catchments in 2007 following consultation with all the local land occupiers concerned. On the south side of the Hurunui all planned work between Maori Gully and Lake Taylor was completed in 2008/09. All trees from large coning conifers (trees of seed producing size) right down to seedlings were removed. This was achieved mainly through 'skid-hopping', with contractors moving from tree to tree by helicopter.

Last year's programme on the north side of the Hurunui focused on land between the Jolly Brook and the Mandamus. This coming year's programme will see a continuation of work in the Mandamus Catchment. The planned control operation is not as straight forward as last year with many of the isolated 'wildings' situated in dense native scrub cover, making ground and aerial access extremely difficult. Other control options are being considered for this work.

If you have wilding conifers on your property, don't wait until they are large and numerous enough to require a costly control programme. Start your control work now. Remember all plants that have become pests started from a few isolated plants and if dealt with at an early stage, their spread could so easily have been halted.

## RABBIT CONTROL: Best practice is cost effective

Whenever rabbit numbers increase throughout the rabbit-prone areas of Canterbury, many farmers use poison programmes to control numbers. However, a poison programme needs to kill 95% or more of the population to be cost efficient.

Affordability is of course a major factor in planning a poison programme, but failing to use poison at the right time has a cost too.

Delay can result in missing the annual seasonal window, increased bait requirements for a bigger rabbit population and an a larger area needing treatment as rabbits spread. Losses due to pasture damage and soil erosion affect productivity and may mean that supplementary feed is needed. All the above are expensive measures and all are avoidable through good planning.



Live rabbits trying to get over a rabbit-proof fence.

Remember that to purchase or apply poison to bait, you must hold a *Controlled Substances License* or an *Approved Handlers Certificate* or be under the direct supervision of a licence holder. Unless you have the above licenses, you are able to purchase Pindone-based products for use in bait stations only.

### THE GOLDEN RULES FOR A COST-EFFECTIVE POISON PROGRAMME.

- **Seasonal timing**  
Best results are achieved from March to August, outside of the breeding season and when natural feed is scarcest.
- **"Rest" your rabbit population**  
Stop shooting a good 3-4 months before poisoning. Wary rabbits don't take bait readily.
- **Work to a rabbit-proof boundary**  
Work to an effective rabbit-proof boundary and concentrate efforts where rabbit numbers are highest.
- **Use good quality bait and apply at the right intervals**  
Use fresh bait in good condition. With 1080, pre feed twice before using poison bait. Feeds should be around a week apart. If more than two weeks apart, a third light feed is best practice. Monitor bait disappearance.
- **Use ample bait when pre-feeding**  
Dominant animals feed first. If too little bait is available, some rabbits will survive. All areas need to be pre-fed at the correct rate so that all individuals can feed.
- **Ensure the correct toxic loading is used**  
It is an offence to alter labelled toxicity and sub-lethal dosing or poison shyness can result.
- **Consider a bait acceptance trial**  
This is especially important if you have a large treatment area or the cost is considerable.
- **Always use reputable contractors**

To summarise, employing best practice methods and getting the timing right are critical in maintaining efficient and cost effective rabbit control.

## Look out for thistles with yellow flowers

Saffron thistle (*Carthamus lanatus*) is an upright thistle that can grow up to a metre tall, with leaves growing out from the centre to form a rosette. Over time, the leaves of the rosette disappear and the stem divides into many branches. The spines on saffron thistle are prominent and very sharp. The flowers appear from about December through to April and seeds germinate in autumn.

The yellow flower is the key to identifying saffron thistle, as most other thistles have pink or purple flowers.

Saffron thistle is a Total Control Pest Programme plant under Canterbury's Regional Pest Management Strategy (2005). Environment Canterbury carries out all control work on such pests.

There are only a small number of known sites in Canterbury. Environment Canterbury's Biosecurity staff at Amberley want to hear from you if you have seen this plant.



The bright yellow flower that distinguishes saffron thistle from most other thistles